

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PUBLIC HEALTH REPORTS.

UNITED STATES.

Statistical reports of States and cities of the United States—Yearly and monthly.

California—Fresno.—Month of November, 1902. Estimated population, 17,000. Total number of deaths, 27, including 4 from tuberculosis.

Stockton.—Month of November, 1902. Census population, 17,506. Total number of deaths, 25, including 3 from tuberculosis.

CONNECTICUT—Hartford.—Month of November, 1902. Estimated population, 83,000. Total number of deaths, 83, including diphtheria, 1; measles, 3; scarlet fever, 1; whooping cough, 2, and 7 from tuberculosis.

ILLINOIS—Bloomington.—Month of November, 1902. Estimated population, 25,000. Total number of deaths, 24, including enteric fever, 1, and 4 from tuberculosis.

Evanston.—Month of October, 1902. Estimated population, 20,344. Total number of deaths, 21, including diphtheria, 1; enteric fever, 1, and 1 from tuberculosis. Month of November, 1902. Total number of deaths, 16, including diphtheria, 2, and 1 from enteric fever.

Kankakee.—Period from November 1 to November 30, 1902. Estimated population, 13,500. Total number of deaths, 17, and 3 from tuberculosis.

Indiana—Evansville.—Month of November, 1902. Estimated population, 65,000. Total number of deaths, 40, including diphtheria, 1; enteric fever, 7, and 6 from tuberculosis.

Muncie.—Month of November, 1902. Estimated population, 22,670. Total number of deaths, 24, including 1 from diphtheria.

MASSACHUSETTS—Holyoke.—Month of November, 1902. Estimated population, 48,065. Total number of deaths, 70, including scarlet fever, 1, and 7 from tuberculosis.

MICHIGÁN.—Reports to the State board of health, Lansing, for the week ended December 6, 1902, from 74 observers, indicate that measles, erysipelas, smallpox, and dysentery were more prevalent and bronchitis, neuralgia, diarrhea, pleuritis, enteric fever, pneumonia, inflammation of bowels, inflammation of kidney, whooping cough, intermittent fever, and cholera morbus were less prevalent. Meningitis was reported present at 6, whooping cough at 19, measles at 27, diphtheria at

2837

42, smallpox at 43, enteric fever at 62, scarlet fever at 79, and phthisis pulmonalis at 215 places.

MINNESOTA—Winona.—Month of November, 1902. Estimated population, 23,000. Total number of deaths, 13, including diphtheria, 1; enteric fever, 1, and 1 from tuberculosis.

NEW JERSEY—Hudson County.—Month of October, 1902. Estimated population, 409,810. Total number of deaths, 521, including diphtheria, 20; enteric fever, 6; scarlet fever, 2; whooping cough, 1, and 70 from tuberculosis.

Passaic.—Four weeks ended December 6, 1902. Census population, 27, 777. Total number of deaths. 27, including 2 from diphtheria.

NEW YORK.—Reports to the State board of health, Albany, for the month of October, 1902, from 156 cities, towns, and villages, having an aggregate estimated population of 7,467,050, show a total of 9,475 deaths, including diphtheria, 215; enteric fever, 225; measles, 17; scarlet fever, 56; smallpox, 4; whooping cough, 61, and 1,027 from tuberculosis.

OHIO—Cleveland.—Month of November, 1902. Estimated population, 400,000. Total number of deaths, 463, including diphtheria, 27; enteric fever, 12; scarlet fever, 5; smallpox, 18, and 31 from tuberculosis.

Zanesville.—Month of November, 1902. Estimated population, 28,000. Total number of deaths, 31, including diphtheria, 3; enteric fever, 1, and 5 from tuberculosis.

TENNESSEE—Chattanooga.—Month of November, 1902. Estimated population, 40,000—white, 27,000; colored, 13,000. Total number of deaths, 23—white, 14; colored, 9—including diphtheria, 3; enteric fever, 1; smallpox, 1, and 5 from tuberculosis.

Nashville.—Month of November, 1902. Estimated population, 81,320—white, 51,128; colored, 30,192. Total number of deaths, 145—white, 67; colored, 78—including diphtheria, 2; enteric fever, 6; whooping cough, 1, and 32 from tuberculosis.

TEXAS—San Antonio.—Month of November, 1902. Estimated population. 63,000. Total number of deaths, 139, including diphtheria, 2, and 28 from tuberculosis, of which 20 were of nonresidents.

UTAH—Salt Lake City.—Month of November, 1902. Estimated population, 75,000. Total number of deaths, 74, including diphtheria, 6; enteric fever, 2, and 2 from tuberculosis.

Inspection of immigrants.

WEEKLY.

Place.	Week ended.	No. of ves- sels.	No. of immigrants.
Baltimore, Md. Do. Boston, Mass. Philadelphia, Pa. Ponce, P. R. Do. San Juan, P. R. Do. Mayaguez, P. R. Do.	Dec. 6 do Nov. 29 Dec. 6	4 3 9 2 2 1 4 2 1	'1, 109 1, 563 736 274 5 1 131 9 2

MONTHLY.

Place.	Month ended.	No of immigrants passed.	No. re-
Sault Ste. Marie, Mich. Astoria, Oreg. New York, N. Y. Philadelphia, Pa. Ponce, P. R. San Juan, P. R. Subports, Porto Rico Eagle Pass, 1 ex. Gaiveston, Tex.	Nov. 30 do	170 67 47, 733	47 0 353
Ponce, P. R. San Juan, P. R.	do	26 183	
Subports, Porto Rico Eagle Pass, 1ex Gaiveston, Tex	do	94 82	6 0